



The Daniel K. Inouye Solar Telescope Captures Ambient Shock Waves Ignited by Magnetic Vortices

Catherine E. Fischer^{1,2}, Friedrich Wöger², Gangadharan Vigeesh³, Peter H. Keys⁴, Oskar Steiner^{3,5}, José Roberto Canivete Cuissa^{5,6}, Shahin Jafarzadeh^{4,7}, Robertus Erdélyi^{8,9,10}, David B. Jess^{4,11}, Mihalis Mathioudakis⁴, Alexandra Tritschler², Damian J. Christian¹¹, Andre Fehlmann^{2,12}, Sarah Jaeggli^{2,12}, and Thomas Rimmele²

¹ European Space Agency (ESA), European Space Astronomy Center (ESAC), Camino Bajo del Castillo s/n, 28692, Villanueva de la Cañada, Madrid, Spain; catherine.fischer@esa.int

² National Solar Observatory, Boulder, 3665 Discovery Drive, 80303 Boulder, USA

³ Kiepenheuer-Institute for Solar Physics (KIS), Georges-Köhler-Allee 401a, 79110 Freiburg, Germany

⁴ Astrophysics Research Center, School of Mathematics and Physics, Queen's University Belfast, Belfast, BT7 1NN, UK

⁵ Istituto ricerche solari Aldo e Cele Daccò (IRSOL), Faculty of Informatics, Università della Svizzera Italiana (USI), Via Patocchi 57, 6605 Locarno, Switzerland

⁶ Center for Theoretical Astrophysics and Cosmology, Institute for Computational Science (ICS), University of Zurich; Winterthurerstrasse 190, 8057 Zürich, Switzerland

⁷ Rosseland Centre for Solar Physics, University of Oslo, P.O. Box 1029, Blindern, 0315 Oslo, Norway

⁸ Solar Physics & Space Plasma Research Center (SP2RC), School of Mathematical and Physical Sciences, University of Sheffield, Hounsfield Road, Sheffield S3 7RH, UK

⁹ Department of Astronomy, Eötvös Loránd University, Pázmány Péter sétány 1/A, H-1112 Budapest, Hungary

¹⁰ Hungarian Solar Physics Foundation, Petőfi tér 3, H-5700 Gyula, Hungary

¹¹ Department of Physics and Astronomy, California State University Northridge, Northridge, CA 91330, USA

¹² National Solar Observatory, 22 Ohi'a Ku Street, Makawao, HI 96768, USA

Received 2025 December 5; revised 2026 August 15; accepted 2026 August 27; published 2026 October 9

Abstract

One of the first high-cadence chromospheric image sequences recorded with the recently commissioned Daniel K. Inouye Solar Telescope (DKIST) has revealed propagating arc-shaped bright fronts of a few hundred kilometers width. These bright fronts maintain their arc shape over up to one minute, travel with apparent horizontal speeds of $\sim 8 \text{ km s}^{-1}$, and originate from chromospheric bright grains seen at the center of the arc. We identify photospheric *G*-band bright points underneath the chromospheric bright grains, which are typically the signature of small-scale magnetic flux concentrations. The *G*-band bright points are found to interact with colocated vortices prior to the appearance of the chromospheric bright fronts. Images synthesized from three-dimensional magnetohydrodynamic simulations show a striking similarity with the discovered phenomenon and reveal that the arc-shaped bright fronts arise from essentially acoustic, weak shock fronts that propagate in the ambient medium, seemingly triggered by the vortex dynamics of the underlying magnetic flux concentration. In this work, we provide the first hints of a hitherto unknown magnetohydrodynamic process by which vortex-driven torsional Alfvén pulses couple to predominantly acoustic waves that steepen into shocks and dissipate their energies. The detected number density and frequency of these events are at least a factor 10 too small to be a major source of solar chromospheric heating but simulations show a seven times higher number density. We therefore expect to discover many more events with tailored narrow band observations.

Unified Astronomy Thesaurus concepts: Solar chromospheric heating (1987); Solar chromosphere (1479); Solar photosphere (1518); Alfvén waves (23); Solar magnetic bright points (1984); Solar physics (1476); Stellar astronomy (1583)

Materials only available in the online version of record: animations

1. Introduction

Past studies of the Sun, currently the only highly resolvable star, have taught us that the discovery and understanding of the physical mechanisms at work in the solar atmosphere require observations of ever higher spatial and temporal resolution. By and by, such high-resolution observations revealed that the solar surface is abundantly occupied by small-scale magnetic flux concentrations at any given time (R. B. Dunn & J. B. Zirker 1973; T. E. Berger & A. M. Title 1996; T. E. Berger et al. 2004; S. Tsuneta et al. 2008; P. H. Keys et al. 2019; see also A. G. de Wijn et al. 2009 or L. Bellot Rubio & D. Orozco

Suárez 2019 for reviews). These magnetic flux concentrations reach superequipartition field strengths of $\approx 0.15 \text{ T}$, presumably as a consequence of the convective collapse instability (H. C. Spruit 1979; S. Nagata et al. 2008; C. E. Fischer et al. 2009). Furthermore, high temporal resolution of image series of the solar atmosphere as seen, for example, in the visible continuum or the *G* band, show a very dynamic solar atmosphere, changing on timescales of minutes and seconds (P. H. Keys et al. 2026). Magnetic field tracers, such as *G*-band bright points (BPs), are seen to be shuffled about by convective plasma motions, leading to a range of transient events. These (BPs) are a reliable indicator of small-scale magnetic flux concentrations (T. E. Berger & A. M. Title 2001), appearing bright due to the interplay between opacity effects, the magnetic field, and the surrounding dynamic plasma (H. C. Spruit 1976; A. M. Title & T. E. Berger 1996; R. J. Rutten et al. 2001; O. Steiner et al. 2001).



Original content from this work may be used under the terms of the [Creative Commons Attribution 4.0 licence](https://creativecommons.org/licenses/by/4.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

More recently, a variety of small-scale processes involving such magnetic elements have been observed, such as (i) dynamic fibrils (V. H. Hansteen et al. 2006) as slow-mode magnetoacoustic shocks propagating in magnetic conduits, (ii) magnetic portals through which low-frequency (<5 mHz) magnetoacoustic waves can propagate into the solar chromosphere (S. M. Jefferies et al. 2006), (iii) vortices in the chromosphere induced and driven by photospheric magnetic elements (S. Wedemeyer-Böhm et al. 2012), (iv) torsional Alfvén pulses running along magnetic flux concentrations from the photosphere to the chromosphere (J. Liu et al. 2019b), or (v) other oscillatory phenomena and jets such as those found by C. E. Fischer et al. (2017) and S. M. Díaz-Castillo et al. (2024), all linking the solar atmosphere across different physical regimes.

Studies of this connectivity between the solar atmospheric layers are highly sought after because they offer insight to the enigmatic heating problem—the fact that the chromosphere and corona display temperatures far exceeding the temperatures in the underlying photosphere. Multidimensional numerical simulations have long predicted that photospheric drivers can excite magnetoacoustic waves and magnetic twist (O. Steiner et al. 1998; V. Fedun et al. 2011; I. N. Kitiashvili et al. 2012; R. Moll et al. 2012; G. Vigeesh et al. 2012; E. Khomenko & P. S. Cally 2019). The waves and twist produced by the near-surface convection can propagate upward in the solar atmosphere and subsequently deposit their energies, possibly by developing into shocks or by other dissipation mechanisms (P. Ulmschneider et al. 1991; S. S. Hasan & W. Kalkofen 1999; B. De Pontieu et al. 2007; R. Erdélyi & V. Fedun 2007; D. B. Jess et al. 2009; S. Wedemeyer-Böhm et al. 2012; D. B. Jess et al. 2023). In fact, acoustic shocks are observed as periodic brightenings in the Ca II H and K lines of the quiet Sun (M. Carlsson & R. F. Stein 1997) and are thought to be the dominant contributor in the heating of the lower chromosphere (M. Carlsson et al. 2019; Q. Noraz et al. 2026). However, A. Fossum & M. Carlsson (2005) came to the conclusion that high-frequency acoustic waves are not sufficient to heat the solar chromosphere. They inferred this from their analysis of the intensity fluctuation in the ultraviolet range of the solar spectrum as observed from space. On the other hand, the ubiquitous magnetic twist and associated torsional Alfvén waves are hard to dissipate, although numerical simulations reveal waves near the boundaries of chromospheric magnetic vortices, which are likely damped by viscous heating (G. Cherry et al. 2025, 2026).

The National Science Foundation’s Daniel K. Inouye Solar Telescope (DKIST; T. R. Rimmele et al. 2020), currently the largest solar telescope in the world, has been designed for exactly this purpose: to study the Sun at unprecedented spatial resolution. DKIST is equipped with advanced adaptive optics technology and a diverse set of instruments for multi-wavelength capabilities and it has now delivered the first science datasets during the currently ongoing Operations Commissioning Phase. In this work, we report on one of DKIST’s first discoveries, namely the presence of small-scale chromospheric wave fronts over the entire field of view (FoV), propagating for several hundred kilometers. They are seen to emanate from bright chromospheric grains that coincide with photospheric BPs, both tracers of the omnipresent small-scale magnetic flux concentrations (S. Jafarzadeh et al. 2013).

With the aid of self-consistent, realistic numerical simulations, which show events very similar to the observed ones, we obtain valuable context for studying the underlying physical properties of these wave fronts, which turn out to be predominantly acoustic, weak shock fronts, triggered by torsional Alfvén pulses. We provide estimates on the occurrence rate and the potential of solar atmospheric heating by such events and point to the broader significance of this novel process.

The outline of the paper is as follows. In Section 2, we describe the DKIST observations and the two sets of simulations including the details of the spectral line synthesis. Section 3 provides a detailed description of the events detected in the observations and comparison with similar events in the simulated data, aimed at a more detailed interpretation of the observations. In Section 4, we discuss the results, including the energy estimates of the detected events and their role in chromospheric heating.

2. Observations and Simulations

2.1. DKIST Observations with VBI

On 2022 May 26, the DKIST observed a quiet Sun region at disk center between 17:46UT and 19:31UT (DKIST Observation ID: pid_1_36). The mean inferred Fried parameter during the extracted observing sequence was 11 cm (at 500 nm wavelength) reaching a value of 15 cm, at its best period. Observations were carried out with the Visible Broadband Imager (VBI; F. Wöger et al. 2021), one of the first-light instruments at the DKIST. The VBI observes the Sun with high-speed $4k \times 4k$ cameras with a frame rate of 30 s^{-1} . The pixel size at the recorded wavelengths corresponds to ~ 0.011 (8 km) on the solar surface, corresponding to Nyquist sampling of DKIST’s theoretical diffraction limit in the *G* band. The instrument acquired images alternating between two visible wavelength filters, with passbands for the *G* band (CH molecular bandhead at around 430 nm) and the Ca II K spectral line (393 nm). The *G*-band filter has a passband width of 0.5 nm full width at half-maximum (FWHM), while the Ca II K filter has 0.1 nm FWHM. These filters provide information about the plasma of the deep photosphere and the upper photosphere/low chromosphere, respectively. The same dataset (pid_1_36) was analyzed with respect to photospheric BPs by P. H. Keys et al. (2026).

The data were dark and flat-field corrected prior to undergoing speckle reconstruction (F. Wöger et al. 2008). The speckle image reconstruction process uses 80 short exposed frames (25 ms exposure time for Ca II K and 1 ms for the *G*-band channel) as input to estimate the Fourier phases and amplitudes of the object using statistical analysis. The result of the processing is a single, high-resolution image calibrated with the best estimate of the system transfer function to achieve a high photometric accuracy in the reconstruction. The resulting cadence for the Ca II K time series and the *G*-band time series after reconstruction is approximately 6 s for both.

The postspeckle reconstructed images were coaligned prior to being destretched (D. B. Jess et al. 2010) by determining local offsets between successive images. This corrects for any residual spatial distortions present as a result of, e.g., atmospheric turbulence, for the duration of the observations, which were not corrected in situ at the telescope (e.g., through




Figure 1. Wave front event Nr. E1 with the G -band BP in the photosphere, the associated Ca II K bright grain in the low chromosphere, and the emitted arc-shaped wave front. (a) Filter image of the VBI G -band channel showing the deep photosphere. The central granule is surrounded at its periphery by BP-like features, which are manifestations of small-scale magnetic flux concentrations. The BP connected to the wave phenomena discussed here is located within the black square. (b) Filter image of the VBI Ca II K channel showing the low chromosphere/upper photosphere. Within the black square, the bright chromospheric grain is partially surrounded by a bright, arc-shaped front traveling away from it. The colored arcs indicate the position of the bright front at subsequent times (see panel (c) for times). (c) Radiative intensity of the Ca II K channel (normalized to the maximum intensity of the Ca II K bright grain) along the radial direction of the travel path of the bright front. Each profile is the mean of profiles taken in 5° steps between the black dashed radial lines indicated in panel (b). The shaded region around the blue curve ($t = 777$ s) displays the minimum and maximum of the intensity for all curves at that time step, shown as a representative example. The position of the bright front, as inferred from the applied Gaussian fit, is marked with triangles, whereby the colors denote the time step of the observation, indicated by labels. The peak intensity at $s \sim 250$ km corresponds to the bright Ca II K grain from where the arc emanates. (d) G -band image recorded approximately 152 s earlier than the G -band snapshot shown in panel (b). (e) Close-up of the region of interest shown in panel (d). Green arrows display the apparent horizontal velocities obtained through a local correlation tracking method (see Section 3.1.1 for details). The velocities have been averaged over a time range of 60 s centered at the time $t = 622$ s. An animation of the maps of panels (a) and (b) is linked to this figure. The animation loops three times and shows a 207 s long sequence, for which the black rectangles and the dashed colored lines have been omitted. A black circle highlights the event location. A pink star indicates the location of the BP at several selected frames of the Ca II K maps and a pink dashed arc denotes the location of the bright front at times $t = 771$ s, $t = 813$ s, and $t = 844$ s.

(An animation of this figure is available in the [online article](#).)

the use of adaptive optics). The final step involved coaligning the G -band and Ca II K images with respect to each other, using long-lived cospatial features, to allow a comparison of features cospatially between the filter images.

A sequence with the best seeing, spanning a FoV of $12.76 \times 12.76 \text{ Mm}^2$ and with a duration of 26 minutes, is selected for the analysis. The data are one of the few quiet Sun disk-center datasets at such a high cadence and with good seeing available so far from DKIST. Although data were simultaneously acquired with the Visible Spectropolarimeter (ViSP; A. G. de Wijn et al. 2022) instrument, our events did not fall within its FoV.

2.2. Simulations and Synthesis

Within this study, we employ two separate simulation data to compare with the DKIST observations. They were obtained using different initial magnetic field configurations and boundary conditions and generated by two fairly different, fully self-consistent, numerical radiation magnetohydrodynamic (MHD) simulation tools, namely, CO5BOLD (B. Freytag et al. 2012) and Bifrost (B. V. Gudiksen et al.

2011). The CO5BOLD simulation snapshots used (ID: d3gt57g44v50fc_high_cad_sec) have a duration of 15 minutes with a cadence of 2 s and with the computational domain spanning $9.6 \text{ Mm} \times 9.6 \text{ Mm} \times 2.8 \text{ Mm}$. The computational cell size is 10 km in each spatial direction. The vertical extent spans from $z \approx -1240$ km to the top boundary at $z \approx 1560$ km, where $z = 0$ corresponds to the mean Rosseland optical depth $\tau_R = 1$. The initial state contained a homogeneous, vertical magnetic field of 50 G magnetic flux density, which was relaxed over 1.5 hr of physical time until the beginning of the analyzed time series. Continuum and Ca II K maps derived from these data were synthesized using the Lightweaver (C. M. J. Osborne & I. Milić 2021) code. The Ca II K spectral line was synthesized over a wavelength range of ± 0.5 nm around the central wavelength of 393.3663 nm. The synthesis was carried out assuming complete frequency redistribution (CRD) but in certain cases, effects of partial frequency redistribution (PRD) are taken into account. Electron densities are calculated under local thermodynamic equilibrium (LTE) conditions. The Ca II K maps are then constructed by applying the VBI transmission profile. The continuum maps are

Table 1

Apparent Horizontal Propagation Speed, FWHM, and Radial Distance of the Bright Front Relative to the Source Location for Observation and Simulation Events

Event Times	Speed (km s ⁻¹)	FWHM (km)	Maximum Position (km)
Obs. Event Nr. E1	9.2 ± 4.7		
$t = 777$ s		215.9 ± 35.3	548.3 ± 17.8
$t = 789$ s		258.1 ± 72.2 ^a	673.1 ± 45.9 ^a
$t = 826$ s		406.4 ± 82.2	950.5 ± 35.7
$t = 838$ s		429.5 ± 66.5	1068.1 ± 43.6
Obs. Event Nr. E2	7.5 ± 4.4		
$t = 722$ s		266.9 ± 10.2	344.4 ± 11.1
$t = 734$ s		279.1 ± 10.2	405.8 ± 18.1
$t = 740$ s		289.7 ± 9.7	465.4 ± 19.8
Obs. Event Nr. E3	6.3 ± 1.1		
$t = 466$ s		182.7 ± 10.1	280.2 ± 2.6
$t = 472$ s		185.7 ± 5.0	325.8 ± 3.2
$t = 479$ s		186.1 ± 14.5	340.0 ± 4.1
$t = 485$ s		103.4 ± 96.4	396.4 ± 3.0
Sim. Event Nr. S1 (CO5BOLD)	7.6 ± 0.8		
$t = 5764$ s		156.6 ± 4.8	372.5 ± 3.3
$t = 5770$ s		182.4 ± 2.0	361.9 ± 1.4
$t = 5776$ s		193.1 ± 7.3	406.5 ± 3.0
$t = 5782$ s		212.8 ± 4.4	464.7 ± 2.8
Sim. Event Nr. S2 (Bifrost)	15.2 ± 1.1		
$t = 4280$ s		186.7 ± 2.8	343.8 ± 4.7
$t = 4290$ s		136.4 ± 4.9	477.1 ± 4.0
$t = 4310$ s		118.2 ± 8.2 ^a	798.0 ± 13.0 ^a
$t = 4320$ s		191.7 ± 8.1	960.6 ± 14.9

Note. The times refer to the times of the Ca II K frames within the full sequence animation (animation linked to Figure B1).

^a 6 profiles used for the error calculation.

synthesized at a wavelength of 450.287 nm. A detailed description of the CO5BOLD simulation tool is given in B. Freytag et al. (2012).

Within this study, we also use the publicly available Bifrost simulation data with identity number ch024031_by200bz005. A detailed description of the Bifrost simulation tool is given in B. V. Gudiksen et al. (2011). The Bifrost data cube has a cadence of 10 s and a range of 24 Mm × 24 Mm × 17 Mm (768 × 768 × 768 grid points) with grid cells of 31.25 km in the horizontal direction. The range in geometrical height is from $z \approx -2.5$ Mm to $z \approx 14.5$ Mm with respect to the solar surface (i.e., optical depth, $\tau = 1$ at 500 nm). In this case, a magnetic field configuration with an average signed strength of 5 G at $\tau_{500} = 1$ is used to represent a coronal hole. At the bottom boundary, a horizontal field of 200 G is applied to the inflows. This setup allows the field strength to slowly increase over time, with interactions between the existing field and the new flux. For the present work we then use a time period of 400 s in which the average unsigned field strength at $\tau_{500} = 1$ is 40 G (frames 420–439). For the synthesis of the spectral lines, the RH code (H. Uitenbroek 2001; T. M. D. Pereira & H. Uitenbroek 2015) was employed.

In both cases, near the solar surface ($\tau_R = 1$), the initially imposed magnetic field is quickly advected to the intergranular lanes by the granular flow. There, the dynamic pressure

concentrates it to dynamical equipartition field strength of several hundred Gauss. Further intensification to kilogauss field strength proceeds by the convective instability in the presence of magnetic fields, a process known as convective collapse (H. C. Spruit 1979). Also in both cases, vortex flows and associated magnetic twists naturally appear and are not deliberately excited. They are produced by the convective dynamics as a result of the self-consistent numerical solution of the MHD equations together with the equations of radiative transfer. More information on the simulations employed and on the differences between the two simulation data can be found in Appendix A.

3. Results

3.1. Observations

3.1.1. Broadband Imagery in the Ca II K and G-band Channels of VBI

We detect propagating fronts of enhanced radiative intensity in the high-cadence Ca II K sequence. Their distinct arc shapes emanate from their source location, where we find one or more steadily evolving chromospheric bright grains with corresponding BPs, in the G-band filter image.

In Figure 1, we present the observation of an example propagating front, denoted as event Nr. E1. The observation times given throughout the text and in the figures and animations are the times given in the full sequence shown in the animation linked to Figure B1.

Figure 1(b) shows a snapshot of a bright arc, visible from the Ca II K filter image series. The bright arc at time $t = 777$ s is delineated by a blue circular segment and its subsequent development in maroon, pink, and lavender corresponding to times $t = 789$ s, $t = 826$ s, and $t = 838$ s, respectively. The bright front starts from the Ca II K bright grain located at $(x, y) = (4200, 1700)$ km at approximately time $t = 759$ s. The full time sequence of the representations shown in panels (a) and (b) can be followed with the animation linked to Figure 1, and an enlarged version of the Ca II K image within the black square is repeated in the top left panel of Figure B1.

The FoV of the G-band filter image shown in Figure 1(a) is dominated by a large granule surrounded by BPs residing in the adjacent intergranular lanes, which give rise to the bright grains seen in the Ca II K filtergrams. An automated feature identification and tracking algorithm (P. J. Crockett et al. 2010) was utilized to isolate BPs within the G-band images. This algorithm isolates local intensity maxima within the intergranular lanes, followed by performing intensity cross sections in different directions at 5° intervals about the barycenter location in order to accurately determine the BP boundary. Measurement of intensity gradients in these cross sections enable the algorithm to discard false detections (e.g., fragmented granules). The tracking phase of the algorithm looks for similar features that persist between multiple frames in the vicinity of the originally detected BP. The location and size of these identified and tracked BPs are then stored for further analysis, where the sequence of locations serve to determine the velocities and paths of tracks of the BPs (P. H. Keys et al. 2011). Features were identified and tracked in the G-band images to determine the relationship between the bright, propagating arcs observed in the Ca II K images and the underlying magnetic field represented by the BPs. The BP shown in the black square of Figure 1(a) is the result of two BPs that merged at $t = 756$ s, after one of them experienced a




Figure 2. Event Nr. E2 of the DKIST VBI examples showing the magnetic BP located within the region of influence of a vortex in the photosphere, its corresponding signature in the low chromosphere, and the associated bright front signature. The figure is in the same format as Figure 1. For panel (e) the velocities have been averaged over a time range of 30 s centered at the time $t = 646$ s. An animation of the images shown in panels (a) and (b) is available and linked to this figure. The animation loops three times and shows a 140 s long sequence, for which the black rectangles and the dashed colored lines have been omitted. A black circle highlights the event location. A pink star indicates the location of the BP at several selected frames of the Ca II K maps and a pink dashed arc denotes the location of the bright front at $t = 698$ s, $t = 716$ s, and $t = 734$ s.

(An animation of this figure is available in the [online article](#).)

rotating motion. The automated BP tracking reveals that this rotating BP first accelerates to a transverse velocity of $1.1 \pm 0.2 \text{ km s}^{-1}$ before entering a 66 s long rotational phase with an angular velocity of $1.1 \pm 0.1 \text{ deg s}^{-1}$. The angular velocity is inferred from the motion of the BP over this time window using the barycentres identified from the BP tracking routine.

The Fourier local correlation tracking (FLCT) code (G. H. Fisher & B. T. Welsch 2008) was applied to the G-band time series to obtain apparent horizontal velocities. The FWHM of the Gaussian employed for weighing of the correlated subimages was set to $\sigma = 20$ pixels. For event Nr. E1 (Figure 1(e)), we averaged the obtained velocities over a time range of 61 s over the rotation phase of the BP. The output of the FLCT code indicates a vortical plasma motion (swirl) during the same time period. This can be seen by visual inspection of the horizontal velocity field that displays a coherent circulating pattern of velocity vectors colocated with the BP (see Figure 1(e)). This rotation of the surrounding plasma takes place approximately two minutes prior to the launch of the wave in the chromosphere. The swirl is likely the underlying cause of the rotational motion of the BP, as is often seen for magnetic flux concentrations due to the ubiquitous presence of small-scale swirling motions within quiet Sun intergranular lanes (J. A. Bonet et al. 2008; J. Liu et al. 2019a; K. Tziotziou et al. 2023).

At close inspection, the bright arc in the Ca II K filtergrams starts as a small, slightly curved brightening at $t = 759$ s on the

left side adjacent to the BP that had a rotational component to its motion, before traveling as a semicircular arc across the large granule. We find a mean horizontal propagation speed of the bright front of $9.2 \pm 4.7 \text{ km s}^{-1}$ (see Table 1 for inferred parameters and associated deviation limits).

The intensity profile of the bright front along its direction of motion can be approximated with a Gaussian curve. The FWHM of this intensity profile is initially around 200 km, and expands in the last time steps to a mean width of 430 km (Figure 1(c)). We clarify here in advance and will show further below that simulations suggest this to be the intensity distribution over a shock-front surface and therefore should not be mistaken with the actual shock structure. It displays a fine structure within the arc with a striation consisting of bright and dark ridges, best visible from time $t = 807$ s onwards, which may point to the onset of a corrugation instability of the shock surface. The analysis of such an instability and also the possible connection with the magnetic structure of the environment is left for future investigations with the availability of spectropolarimetric data. At $t = 856$ s, the arc has traveled to a distance corresponding to approximately the middle of the underlying photospheric granule. At this time, several brightenings can be distinguished within and around the arc before it diffuses and becomes untraceable.

Table 1 lists the apparent horizontal propagation speed, FWHM, and radial distance of the bright front relative to the source location for observation and simulation events. We select seven radial directions from the chromospheric bright




Figure 3. Event Nr. E3 of the DKIST VBI examples showing the magnetic BP located within the region of influence of a vortex in the photosphere, its corresponding signature in the low chromosphere, and the associated bright front signature. This figure is in the same format as Figure 1. For panel (e) the velocities are shown for the time $t = 390$ s. An animation of the images shown in panels (a) and (b) is available and linked to this figure. The animation loops three times and shows a 122 s long sequence, for which the black rectangles and the dashed colored lines have been omitted. A black circle highlights the event location. A pink star indicates the location of the BP at several selected frames of the Ca II K maps and a pink dashed arc denotes the location of the bright front at $t = 503$ s, $t = 521$ s, and $t = 539$ s. (An animation of this figure is available in the [online article](#).)

grain to the arc-shaped bright front, separated by 5° intervals for observations event Nr. E1 for determining intensity profiles. For events Nr. E2 and E3, we also choose seven radial directions with intervals of 1° with the exception of the first and last interval of event Nr. E2 with 2° and 12° , respectively. The radial intensity profiles are fitted using the `lmfit` python routines and employing a Gaussian fit to the radial profile of the bright front. The front position (maximum of the Gaussian fit) and the FWHM are determined by fitting the mean of the seven profiles. The errors are the standard deviation of the fitting results obtained from the individual intensity profiles. We compute the speed of the bright front from the maxima found with the Gaussian fit to the mean profile for the time steps listed in Table 1. The average speed is the mean of these speeds whereby the error is the standard deviation of the individual speeds obtained for the mean profiles. The same method was applied to the simulation data presented in Section 3.2.1 with the radial profiles taken at 1° intervals for simulation event Nr. S1 as well as for simulation event Nr. S2.

3.1.2. Additional Similar Events in the Observation Data

Nine more events exhibiting an evolution similar to event Nr. E1 are detected within the examined image sequence. This implies a rate of 0.14 events per Mm^2 per hour within the sequence. These examples are not as prominent as the example seen in event Nr. E1 and decay faster. Sample images for all 10 events are shown in the Appendix B as Figure B1, while

results for the apparent horizontal propagation speed, FWHM, and position of the wave front are provided in Table 1 for event Nrs. E1–E3 for which these quantities are best identifiable.

Events Nr. E2 and E3 are presented in the same manner as in Figure 1, including their animation in Figures 2 and 3, respectively. They also show in the Ca II K filter images an arc-shaped enhanced wave front originating from a Ca II K bright grain that coincides with one or several photospheric BPs. Figure 2 (event Nr. E2) shows the time instance of an event in which a chain of BPs trace the advecting swirling motion (see G-band images of the full time sequence in the animation linked to Figure 2 and the flow field obtained from local correlation tracking (LCT) in Figure 2(e)). The Ca II K bright front launched at approximately $t = 691$ s, a minute after the appearance of the swirl, is much fainter than in event Nr. E1, yet is clearly visible. We are able to trace the front, of width 266.8 ± 10.2 km, for about 50 s in which time span it propagates at an apparent horizontal speed of $7.5 \pm 4.4 \text{ km s}^{-1}$ before becoming less discernible when merging with surrounding brightenings. Event Nr. E3 shows a BP caught in a vortex and assuming a “U”-shaped form (see Figure 3 and accompanying animation). The evolution of all 10 events is shown in the animation of Figure B1. Although the events exhibit a similar overall behavior, we see a variety in visibility, lifetime, propagation speed, and even shape of the bright front with some being close to circular, and others becoming visible in different locations along the arc as time




Figure 4. Comparison of the observational case Nr. E1 with a simulated atmosphere and synthesized blue continuum and Ca II K maps. (a) Volume rendering of the temperature above 5000 K of simulation event Nr. S1 (at $t = 5814$ s) with blue magnetic field lines representing the twisted magnetic vortex structure. The bright front appears as a thin, arc-shaped temperature enhancement on one side of the vortex illustrated by the dashed pink semicircle with arrows indicating the direction of propagation. The shock front proper is visible as the sheet of enhanced temperature slightly inclined with respect to a horizontal plane. The location of the line of sight used in Figures 7 and D2 at $(x, y) = (3435 \text{ km}, 3755 \text{ km})$ is shown as a white line on the right-hand side in the 3D view. (b) DKIST/VBI observations of event Nr. E1 at time $t = 826$ s. The pink dashed arc shown in the Ca II K image represents the location of the bright front, while arrowheads indicate the direction of travel of the front. (c) The top map shows the line-core Ca II K map referring to the low chromosphere/upper photosphere and the bottom map shows the continuum intensity at 450 nm referring to the nearly same photospheric region as the observed G-band image, both of which were computed from the numerical simulation data. The middle map displays the line integral convolution of the horizontal velocity field at $z = 750$ km with the colors denoting the vertical velocity, v_z , where red indicates downflows and blue indicates upflows. An animation of panel (a) is linked to this Figure. The time sequence spans a duration of 68 s and shows the evolution of the temperature and magnetic lines of force. In the time sequence, the bright front appears as a thin, arc-shaped temperature enhancement on one side of the vortex at the top boundary of the subdomain. The shock front appears as an extended sheet of enhanced temperature, slightly inclined with respect to the horizontal plane.

(An animation of this figure is available in the [online article](#).)

progresses. We will return to this observation in Section 3.2.2, where we speculate on the possible cause of these differences.

For events Nr. E1 and E3 (Figures 1 and 3), the BP is close to the swirl center—this configuration very likely excites a torsional Alfvén pulse. The time delay between the appearance of the photospheric swirl to the launch of the bright arc seen in Ca II K is ~ 130 s for event Nr. E1 and ~ 70 s for event Nr. E3. For event Nr. E2 the situation is different. Rather, we find there a configuration where a chain of small G-band BPs move relative to a single BP, both being advected by the velocity field in the periphery of a vortex. That relative motion is likely

sufficient to excite a torsional wave or pulse but we cannot exclude that other modes, such as sausage, kink, and/or flute, are excited as well. In fact, such relative motion of G-band BPs have been observed to be the photospheric basis of chromospheric swirls (S. Wedemeyer-Böhm & L. Rouppe van der Voort 2009), which are typically a signature of torsional Alfvén pulses (A. F. Battaglia et al. 2021). For observation case Nr. E2, the time delay between the appearance of the photospheric swirl to the launch of the bright arc seen in Ca II K is ~ 45 s. We observe in 7 out of 10 cases a vortical motion in the photosphere prior to the emission of the bright




Figure 5. Simulation event Nr. S1 in the same format as Figure 1. (a) Synthesized continuum map at the wavelength 450.287 nm. The photospheric bright continuum point at $(x, y) = (1000 \text{ km}, 1000 \text{ km})$ is located at the source of the chromospheric arc-shaped front seen in panel (b). (b) Synthetic Ca II K map computed under PRD. The colored arcs indicate the position of the bright front at subsequent times (see panel (c) for times). (c) Normalized intensity of the Ca II K map in panel (b) along the radial direction of the travel path of the bright front. Each profile is the mean of profiles taken in 1° steps between the black dashed radial lines indicated in panel (b). The position of the bright front is marked with triangles, whereby the colors denote the time step indicated by labels. The procedure to obtain these values is equivalent to the procedure used for the observational analysis. (d) Continuum image computed 40 s earlier than the Ca II K snapshot shown in panel (b). (e) Close-up of the region of interest shown in panel (d). Green arrows display the horizontal velocities extracted directly from the simulation at height $z = 0 \text{ km}$. For panel (e), the velocities are shown for the time $t = 5750 \text{ s}$. An animation of the continuum and Ca II K maps shown in panels (a) and (b) is available and linked to this figure. In the time sequence of duration 82 s, the bright arc-shaped front is marked in the Ca II K map at times $t = 5800 \text{ s}$, $t = 5808 \text{ s}$, and $t = 5818 \text{ s}$ by a dashed pink arc in the replay. The pink star indicates the location of the photospheric BP during the time span of the event. (An animation of this figure is available in the [online article](#).)

front—in three cases a swirl is not easily identified. The temporal distance between the occurrence of the vortex and the bright front is estimated by the time difference between the full development of the swirl in the horizontal velocity field as derived with correlation tracking from the *G*-band image sequence and the first signature of the bright front as detected by visual inspection of the Ca II K image sequence. The time delay for all cases is between about 1 and 2 minutes, which approximately corresponds to the distance between photosphere and low chromosphere when adopting a reasonable characteristic speed of 10 km s^{-1} . Therefore, we consider the connection between vortical flow and launch of the bright front as very probable, although we cannot prove it from the observational data alone.

3.2. Simulations

3.2.1. Bright Front in Numerical Simulations

In the following, we concentrate on simulation event Nr. S1 from the CO5BOLD data and simulation event Nr. S2 from the Bifrost data. For simulation event S1, synthetic Ca II K maps have been generated by applying the Lightweaver synthesis code (C. M. J. Osborne & I. Milić 2021) to the full time series. Figure 4(c) shows a map of the Ca II K intensity in the line core as computed under partial redistribution (PRD), while the right panel of the animation linked to Figure 5 shows the

sequence of maps of the Ca II K line computed under complete redistribution (CRD) and convolved with the VBI transmission profile using the approximation given in F. Wöger (2014). The Ca II K maps of simulation event S2 were produced with the RH code (H. Uitenbroek 2001; T. M. D. Pereira & H. Uitenbroek 2015) under CRD, again applying the VBI filter profile, and are shown in the right panel of the animation linked to Figure 6. Both simulation sequences show wave signatures remarkably similar to those found in the observations, in particular, arcs of enhanced Ca II K emission originating from magnetic flux concentrations and showing an apparent propagation in the horizontal direction. We thus encounter the same phenomenon while utilizing simulation data from two different simulation tools with different initial magnetic configurations and boundary conditions, which testifies to the robustness of the process. The continuum maps of simulation event Nr. S1 reveal a BP at the source location of the front. We also produced synthetic maps of the detailed *G*-band channel, which look very similar to the maps of the continuum at 450.287 nm, as expected. In particular, the continuum BP is equally well visible as a *G*-band BP. Appendix C gives the synthetic imagery of an additional simulation event, Nr. S3, found in the simulation data.

Figure 5 shows the bright front and its radial profiles in the same format as the observations. A photospheric swirl is located cospatial with the magnetic flux concentration as seen




Figure 6. Simulation event Nr. S2 in the same format as Figure 1. (a) Synthesized continuum map at the wavelength 450.287 nm. The photospheric bright continuum point at $(x, y) = (2300 \text{ km}, 2600 \text{ km})$ is located at the source of the chromospheric arc-shaped front seen in panel (b). (b) Synthetic Ca II K map computed under CRD. The colored arcs indicate the position of the bright front at subsequent times (see panel (c) for times). (c) Normalized intensity of the Ca II K map in panel (b) along the radial direction of the travel path of the bright front. Each profile is the mean of profiles taken in 1° steps between the black dashed radial lines indicated in panel (b). The position of the bright front is marked with triangles, whereby the colors denote the time step indicated by labels. The procedure to obtain these values is equivalent to the procedure used for the observational analysis. (d) Continuum image computed 80 s earlier than the Ca II K snapshot shown in panel (b). (e) Close-up of the region of interest shown in panel (d). Green arrows display the horizontal velocities extracted directly from the simulation at height $z = 0 \text{ km}$. For panel (e), the velocities are shown for the time $t = 4210 \text{ s}$. An animation of the continuum and Ca II K maps shown in panels (a) and (b) is available and linked to this figure. In the time sequence of duration 390 s, the bright arc-shaped front is marked in the Ca II K map at times $t = 4290 \text{ s}$ and $t = 4300 \text{ s}$ by a dashed pink arc in the replay. The pink star indicates the location of the photospheric BP during the time span of the event. (An animation of this figure is available in the [online article](#).)

in Figures 5 (d) and (e). The mean velocity of the front as derived from the synthetic Ca II K images is $7.6 \pm 0.8 \text{ km s}^{-1}$. The width of the front increases from $156.6 \pm 4.8 \text{ km}$ to $212.8 \pm 4.4 \text{ km}$ in 18 s. Simulation event Nr. S2 also shows a continuum BP at the source of its traveling bright front (see Figure 6). The front exhibits a faster apparent horizontal propagation of $15.2 \pm 1.1 \text{ km s}^{-1}$ with the width of the bright front changing only marginally during the sequence. These numbers are comparable to the corresponding numbers seen in observations within their range in variation.

3.2.2. Physical Topography of the Bright Front

The simulation data offer the possibility to study the bright front in more detail. Simulation event Nr. S1 is showcased in Figure 4. The BP at the source location of the wave (Figure 5 and animation) exhibits a magnetic field strength of around 0.05 T (see also the animation linked to Figure 4). Tracing the magnetic field lines (blue lines in Figure 4(a)) reveals that they are twisted, implying the presence of a magnetic vortex (S. S. A. Silva et al. 2021). Examining the horizontal velocities at different heights of the model, we find that the twisted magnetic flux tube is located within a plasma swirl, present throughout the photosphere, and reaching the chromosphere. The swirl can be found in the middle image in Figure 4(c), which shows the line integral convolution of the horizontal velocity field for a specific height in the model atmosphere

(750 km) superimposed with the vertical velocity, v_z , with blue colors denoting flows in the upward direction and areas in red denoting flows in the downward direction. From the animation linked to Figure 4, one recognizes a temperature enhanced wave front (labeled “Shock front” in Figure 4(a)) in the surrounding, ambient medium of the twisted magnetic flux concentration. The front is slightly inclined with respect to a horizontal plane, moving upward, and steepening into a hot, sharp front. Thus, from the simulation we conclude that it is a vertically propagating, inclined shock front that causes the observed apparent horizontally moving bright front, which in fact is the shock-front surface seen face-on. The vertical plasma motions around the shock front show downward flows upstream of the shock front and upward flows behind it. At this point, we cannot compare this finding with observations for lack of information from spectral data but shall become a topic of future investigations.

Figure 7 shows the temperature and Ca II K intensity as a function of time along the line of sight at pixel location $(x, y) = (3435 \text{ km}, 3755 \text{ km})$, indicated by the white vertical line in Figure 4(a). There, we identify the shock wave as a fast magnetosonic, predominantly acoustic wave in a high plasma- β ($\beta > 1$) environment (black arrow in Figure 7), where plasma- β is the ratio of gas pressure to magnetic pressure. The plasma- β is approximately equal to the ratio of sound speed to Alfvén speed (c_s/v_A). The contours of $\beta = 10$ and $\beta = 0.1$ in




Figure 7. Time–distance diagram of the temperature from simulation event Nr. S1. The diagram shows the passage of the shock front and the corresponding variation in Ca II K filter intensity at location $(x, y) = (3435 \text{ km}, 3755 \text{ km})$ indicated by the vertical white line in Figure 4(a). The top panel shows the variation in the filter intensity (normalized to the intensity integrated over the full line width) calculated assuming CRD (black curve) and when taking PRD (red curve) into account. The bottom panel shows the temperature. The black arrow points at the temperature enhanced traveling shock front, which splits into two branches when crossing the $\beta = 1$ surface (black curve). The gray dashed contours indicate the layers of $\beta = 0.1$ and 10. The white line indicates the Rosseland optical depth $\tau_R = 1$. The dotted paths show the trajectory of a test particle moving with the fast speed $\sqrt{c_s^2 + v_a^2}$ plus the line-of-sight component of the velocity (blue) and the sound speed (c_s) plus the line-of-sight component of the velocity (white), while v_a is the Alfvén speed. Up to the branching, the two trajectories stay very close to each other because the Alfvén speed is low, and therefore the wave is a fast and predominantly acoustic wave.

Figure 7 indicate the region where the plasma is high- and low- β , respectively. In the case of the high- β regime (i.e., $\beta > 1$), corresponding to $c_s/v_A > 1$), the sound speed is greater than the Alfvén speed, and therefore the wave that follows the white dotted trajectory is mainly driven by gas pressure perturbation running at the fast (acoustic) speed and is consequently considered to be a fast, predominantly acoustic wave. We clearly see the temperature enhancement associated with this fast (predominantly acoustic) wave in the high- β region. As this wave crosses the $\beta = 1$ surface into the low- β regime where the Alfvén speed is larger than the sound speed, the wave splits into two branches: a fast wave, predominantly magnetic in nature, follows the blue dotted trajectory and a slow wave, predominantly acoustic in nature, closely follows the white dotted sound speed trajectory. More details of the shock wave identification are given in Appendix D and corresponding Figures D1 and D2. The top panel in Figure 7 captures the change in the Ca II K filter intensity at the corresponding location as the shock front traverses the line-forming region.

As the inclined shock front moves into the height range of the Ca II K line formation, it closely mimics the observation by forming an arc-shaped front traveling in the horizontal direction as a consequence of projection effects. This also explains why the radiative intensity of the bright front may increase with time (as it does in observation case Nr. E1) and not monotonically decrease as one might expect from a dissipating shock front. Both observation and simulation

events show a variation in the width and appearance of the bright fronts seen in the Ca II K diagnostics (see also Appendix C). The interplay between shock propagation direction, projection effect, and spectral response is likely responsible for this variety.

3.2.3. Torsional Alfvén Pulse

The simulation data show propagating swirls, which can be traced from the photosphere to the chromosphere, where they appear in close vicinity to the bright front. Figure 8 shows the horizontal component of the velocity field for simulation event Nr. S1 in the left column of panels and for simulation event Nr. S2 in the right column. For both events, a swirl at the chromospheric level (top row) is seen in the location from where the bright front emanated. It is preceded by a swirl at the photospheric level (bottom row). The time delay between the photospheric swirl (shown in Figure 8(b) for the simulation event Nr. S1) and the first appearance of the bright front in the synthetic Ca II K maps at time $t = 5780 \text{ s}$ (visible from the animation linked to Figure 5) is 96 s for simulation case Nr. S1 and 50 s for simulation case Nr. S2. For the case of simulation event Nr. S1, Figure 9 shows the swirling strength and the magnetic swirling strength as a function of height and time. Here, the swirling strength and magnetic swirling strength are defined according to A. F. Battaglia et al. (2021); they can be considered to be a measure of vortical motion and magnetic twist, respectively. From this, one can see the swirling motion propagating along the green dotted curve from the upper




Figure 8. Horizontal velocities from the simulated atmosphere. (a) The temperature above 5000 K is shown as a background image to the horizontal velocity depicted with blue arrows for event Nr. S1 (CO5BOLD) at a height of $z = 750$ km at time $t = 5814$ s. (b) Continuum intensity at a wavelength of 450 nm as a background to the horizontal velocity for event Nr. S1 at a height of $z = 0$ km at time $t = 5684$ s. (c) The temperature above 5000 K is shown as a background image to the horizontal velocity depicted with blue arrows for event Nr. S2 (Bifrost) at a height of $z = 1200$ km at time $t = 4310$ s. (d) Continuum intensity at 450 nm as a background to the horizontal velocity for event Nr. S2 at a height of $z = 0$ km at time $t = 4210$ s.

photosphere to the low chromosphere in phase with a propagating twist in the magnetic field, where the magnetic field points opposite to the propagation direction. This phase relation and polarity is indicative of a propagating torsional Alfvén pulse in this region. More explanations are provided in the caption to Figure 9.

4. Discussion

In this paper, we report on the first DKIST VBI observation revealing chromospheric waves in the form of bright fronts in the spectral passband of Ca II K that expose their source by manifesting a distinct arc shape. We detect at their apparent origin chromospheric bright grains and colocated *G*-band BPs, the typical signatures of photospheric magnetic flux concentrations. Furthermore, we find atmospheric vortices at the

location of the *G*-band BPs prior to the launch of the chromospheric waves. Previous observational studies have inferred ubiquitous propagating Alfvén waves or pulses from spectroscopic observations (D. B. Jess et al. 2009; J. Liu et al. 2019b). However, these have not been reported to be connected to chromospheric traveling fronts as detected in our current study. Previously, a case of a radially traveling front has been reported much higher up in the transition region following a small-scale jet triggered within a magnetic element (E. Scullion et al. 2011). However, this event was not associated with a vortex as in our study. S. H. Park et al. (2016) found propagating arcs around a vortex region, which they said would become the subject of future work. No further details are given in the paper and it is not clear if this is the same phenomenon as studied in the present paper. Also, previous analysis of simulation data have shown that torsional




Figure 9. Time–distance diagrams of swirling strength and magnetic swirling strength. The top and bottom panels show the vertical component of the swirling vector λ_z , and the magnetic swirling vector λ_z^B , respectively, as a function of height and time for simulation event Nr. S1 shown in Figure 4. Swirling vector and magnetic swirling vector can be considered a generalization of fluid vorticity and magnetic twist, respectively (see A. F. Battaglia et al. 2021). The quantities are averaged over a horizontal area of $300 \times 300 \text{ km}^2$ centered on the event at coordinates $(x, y) = (3195, 3195) \text{ km}$. The dotted paths show for comparison the trajectory of a test particle moving with the Alfvén speed plus vertical component of velocity. The speeds are averaged over the horizontal area. The light shaded part above the black curve shows the region where the average magnetic field is pointing upward; below it, it is pointing downwards. The ridges along the dotted curves have the same sign in λ_z and λ_z^B and move approximately with the Alfvén speed, indicative of a torsional Alfvén pulse that travels along the flux concentration with oppositely directed magnetic field. The ridges start to become visible in the top of the photosphere only where the magnetic field starts to become dynamically relevant. Also, their appearance is sensitive to the precise location of the chosen line of sight and area due to the horizontal displacement of the vortex and magnetic field with time.

Alfvén pulses that propagate from the photosphere into the chromosphere very commonly occur (A. F. Battaglia et al. 2021; J. R. Canivete Cuissa & O. Steiner 2024) but again without the detection of a horizontally traveling bright front.

The vortices detected in our study are capable of inducing torsional motions within the magnetic flux concentrations. Due to the magnetic flux concentrations exhibiting an inhomogeneous atmosphere, in particular, torsional Alfvén pulses can couple to other types of MHD waves (D. S. Bloomfield et al. 2007; J. Scalisi et al. 2021). In particular, we find coupling to external, nonlinear, predominantly acoustic waves in the ambient atmosphere that may steepen into shock fronts, such as the one seen in Figure 4(a). We note that the events bear a similarity to penumbral running waves identified as “in effect low- β slow-mode waves propagating along the magnetic field” (D. S. Bloomfield et al. 2007). Although we do see a slow-mode (field-aligned) component in the low- β plasma region in simulation event S1 (above the black $\beta = 1$ contour in Figure 7), the shock front already appears in the high- β region at about 5000–6000 K (below the black $\beta = 1$ contour in Figure 7), hence stemming from a fast, predominantly acoustic wave.

Furthermore, while previous studies (D. B. Jess et al. 2009; J. Liu et al. 2019b) show that ubiquitous torsional Alfvén pulses are present in the solar atmosphere, we now provide new evidence of vortical structures emitting magnetosonic waves that shock and therefore dissipate in their vicinity. This is a major discovery of the present study and is of general astrophysical interest since it points us to an unexpected route of dissipating torsional Alfvén waves, which otherwise would hardly dissipate. It may also spur MHD laboratory experiments similar to those of F. Stefani et al. (2021) and is reminiscent of MHD technologies of hypersonic applications.

4.1. Energy Estimates

This study also naturally raises the question of how energetic these events are. We estimate for simulation event Nr. S1 that the shock event supplies an average power of $\sim 2.46 \times 10^{15} \text{ W}$ (see Appendix D for further details). The required energy flux to sustain the lower chromosphere is estimated at $\sim 2 \times 10^{15} \text{ W Mm}^{-2}$ (see E. Priest 2014, Table 1.1 on page 8). Assuming that a typical shock event provides the power found in the simulation and all of its energy is used for replenishing chromospheric radiative losses, then 0.8 per

Mm^2 events would be required to be constantly present to energize the lower chromosphere. So far, we found a much lower occurrence rate of 0.14 events per Mm^2 per hour. Torsional Alfvén pulses have been observed to occur with a rate of up to 0.08 per Mm^2 at any time (I. Dakanalis et al. 2022) and simulations suggest a density of up to $\sim 1 \text{ Mm}^{-2}$ at any time (J. R. Canivete Cuissa & O. Steiner 2024), which indicates the potential of a much higher occurrence rate for swirl driven shock waves. However, taking the present results at face value, these events remain a contributor but not the main contributor to chromospheric heating.

4.2. Contribution to the Solar and Stellar Chromospheric Basal Flux

In a broader context, more than 85% of the stars in our galaxy have surface hydrogen convective envelopes, such as our own Sun, and exhibit solar-like activity and variability. There is increasing evidence that solar-like BPs (B. Beeck et al. 2015; R. G. Salhab et al. 2018) exist in these objects while magnetoacoustic waves are thought to provide a basal energy source for the heating of their chromospheres (M. I. Pérez Martínez et al. 2011). To provide the solar chromospheric H and K basal flux, an occurrence rate of 0.14 per Mm^2 at any time would suffice (see Appendix D for further details). This possibility is of particular interest for stellar physics with regard to the relative chromospheric H and K basal emission index as a function of color index (S. Boro Saikia et al. 2018; J. R. Canivete Cuissa et al. 2024).

4.3. Future Direction

The observed events represent one of the rare observations of how the readily available kinetic energy present in the photosphere can be channeled into the outer regions of the solar atmosphere. Other diagnostics may reveal events similar to the ones presented in this paper but occurring in different layers of the solar atmosphere, possibly even in the corona. In consequence, they warrant further analysis, such as a multi-wavelength spectropolarimetric study, to reveal their thermodynamic and magnetic structure. Identifying the possible connectivity and modes of energy conversion throughout the solar atmosphere will lead to advances in deciphering the signal of unresolved stars. While we only catch glimpses of the processes and phenomena that arise from the surface of other stellar photospheres (X. Dumusque et al. 2011; B. S. Lakeland et al. 2024), the DKIST observations reported here allow us to investigate the complex interplay between the plasma and the magnetic field at an unprecedented level of detail.

Acknowledgments

The research reported herein is based on data collected with the Daniel K. Inouye Solar Telescope (DKIST), a facility of the National Solar Observatory (NSO). NSO is managed by the Association of Universities for Research in Astronomy, Inc., and it is funded by the National Science Foundation. Any opinions, findings, and conclusions or recommendations expressed in this publication are those of the authors and do not necessarily reflect the views of the National Science Foundation or the Association of Universities for Research in Astronomy, Inc. DKIST is located on land of spiritual and cultural significance to Native Hawaiian people. The use of

this important site to further scientific knowledge is done so with appreciation and respect.

The authors wish to acknowledge scientific discussions with the Waves in the Lower Solar Atmosphere (WaLSA; <https://www.WaLSA.team>) team, which has been supported by the Research Council of Norway (project no. 262622) and The Royal Society (award no. Hooke18b/SCTM). This research was also supported by the International Space Science Institute (ISSI) in Bern, through ISSI International Team project 502 (WaLSA: Waves in the Lower Solar Atmosphere at High Resolution).

J.R.C.C. and O.S. acknowledge support by the Swiss National Science Foundation under grant IDs 200020_182094 and 200020_15103 and the University of Zürich under grant UZH Candoc 2022 ID 7104.

O.S. acknowledges HPC resources by the Swiss National Supercomputing Center (CSCS) under project ID s560.

R.E. is grateful to STFC (UK, grant number ST/M000826/1) and acknowledges support from NKFIH Hungary (OTKA, grant No. K142987 and Excellence Grant, grant nr TKP2021-NKTA-64) and NSF China (PIFI, grant No. 2024PVA0043).

S.J. and D.B.J. wish to thank the UK Science and Technology Facilities Council (STFC) for the consolidated grants ST/T00021X/1 and ST/X000923/1. D.B.J. also acknowledges support from the UK Space Agency for a National Space Technology Programme (NSTP) Technology for Space Science award (SSc 009), as well as The Leverhulme Trust for grant RPG-2019-371.

M.M. acknowledges support from the Science and Technology Facilities Council (STFC) under grant No. ST/P000304/1, ST/T001437/1, ST/T00021X/1 and ST/X000923/1, as well as The Leverhulme Trust for grant RPG-2019-371.

Ivan Milic is gratefully acknowledged for his valuable assistance with the Lightweaver code.

Appendix A

Differences Between the Two Simulation Data

In-depth analysis of simulation and synthetic data has been primarily carried out on the CO5BOLD data. However, we also use a Bifrost simulation and synthetic data to consolidate the analysis. CO5BOLD data are used in the first place for their higher spatial resolution (10 km versus 31.25 km computational cell size) and lower diffusivities (see further below) compared to the available Bifrost data. The initial condition for the magnetic field of the two models is fundamentally different. In both cases, the initial condition for the magnetic field remains imprinted on the subsequent time evolution. CO5BOLD data started with a homogeneous vertical field with a strength of 50 G, mimicking a predominantly unipolar magnetic network patch, while Bifrost data started with a unipolar vertical field of 5 G, mimicking the mean signed field strength of a coronal hole, and was subsequently supplied with horizontal magnetic field with a strength of 200 G that slowly entered across the bottom boundary in regions of upflows, leading to an averaged unsigned magnetic field strength in the photosphere of 40 G. This situation results in a more turbulent, mixed polarity magnetic field at the solar surface than in the case of the CO5BOLD data. Details on the boundary conditions are given in F. Calvo (2018) (Section 2 of the PhD Thesis) for the CO5BOLD data and in M. Carlsson et al. (2016) and under <https://sdc.uio.no/search/simulations> for the Bifrost data and

in particular in A. J. Finley et al. (2022) for the Bifrost model ch024031_by200bz005.

The two datasets also differ regarding diffusivities. The CO5BOLD model features a shock capturing Riemann type numerical solver (HLL; A. Harten et al. 1983) based on a reconstruction scheme (B. Freytag 2013) that keeps diffusivities locally minimal while preserving the stability of the numerical solution. This renders, in particular, relatively sharp shock fronts. Bifrost, on the other hand, uses an explicit global and local-specific viscosity and resistivity, large enough to keep the solution stable in time. For the CO5BOLD data, we estimate an effective kinematic viscosity and plasma resistivity near the top of the photosphere of $\nu_{\text{eff}} = 4.0 \times 10^{10} \text{ cm}^2 \text{ s}^{-1}$ and $\eta_{\text{eff}} = 3.3 \times 10^{10} \text{ cm}^2 \text{ s}^{-1}$, respectively (F. Riva & O. Steiner 2022). For the Bifrost data, no such estimate is available but diffusivities are likely larger mainly, but not

only, as a result of the lower spatial resolution. The Bifrost data includes a corona with Spitzer heat conductivity and a transition region to the chromosphere with optically thin radiative cooling (but H-ionization is treated in LTE). None of this physics is included in the CO5BOLD data, which reaches to a height of 1560 km above optical depth $\tau_{\text{R}} = 1$ only. Despite these significant differences in the production of the simulation data, we find in both the CO5BOLD and the Bifrost data events that much resemble the observed ones corroborating that these arise from a robust process not depending on the details of the initial data and boundary conditions.

Appendix B Additional Observation Examples

The following figure (Figure B1) shows sample images for all 10 events detected in the observational dataset.




Figure B1. Collage of observed bright front events. The panels show time instances of the events seen in Ca II K with tick marks set 200 km apart. Part of the bright arc-shaped front is marked by a dashed pink arc. A pink star indicates the location of the Ca II K bright grain and corresponding G-band BPs from where the bright front emanates. An animation showing the temporal sequence of all events is linked to the present figure. The animation shows the continuum maps in the left panel and the Ca II K maps in the right panel. In the time sequence of duration 1541 s, each of the 10 events is highlighted by a colored circle during its occurrence and numbered according to the event numbers given in this figure.

(An animation of this figure is available in the [online article](#).)

Appendix C

Additional Similar Events in the Simulation Data

We identify through visual inspection of a 632 s long CO5BOLD high-cadence sequence 16 events showing a scenario similar to the above-described cases Nrs. S1 and S2, which all are associated with a swirl as well as with a photospheric magnetic flux concentration. We find, similar to the observations, apart from the basic common appearance, a variety in duration, shape, and lifetime of the bright fronts. Figure C1 shows a time instance and corresponding animation of simulation event Nr. S3 from the CO5BOLD data. The animation linked to Figure C1 depicts, as an example, an event

with a wave front that does not exhibit a large apparent horizontal propagation speed. We found one more event with a shock-front topography very similar to simulation case Nr. S1; in the remaining cases, the unambiguous tracking of individual shock fronts was challenging due to the presence of multiple temperature peaks and wave activity from various other sources. The number of found events is, as in the observations, most likely a lower limit for these types of events due to the likelihood of events going unnoticed during visual inspection or forming in a different height range. The occurrence rate is by a factor of about seven larger in the simulations compared to the observations which are indeed susceptible to data quality and seeing limitations.




Figure C1. Synthesized maps from simulation event Nr. S3: The animation shows in the right panel the synthesized Ca II K passband filter maps referring to the low chromosphere/upper photosphere and the left panel maps of the continuum intensity at 450 nm as a close proxy for the observational *G*-band images, both computed from the CO5BOLD numerical simulation. The time sequence of duration 102 s shows the bright arc-shaped front marked in the Ca II K map at times $t = 5896$ s, $t = 5904$ s, and $t = 5912$ s by a dashed pink arc in the replay. The pink star indicates the location of the photospheric BP during the time span of the event. (An animation of this figure is available in the [online article](#).)

Appendix D

Wave/Shock Identification and Energy Estimate

For simulation event S1, the shock front is identified as a sharp temperature enhanced front in the computational domain (see Figures 4 and 7). We then construct a surface in the three-dimensional space based on this peak in temperature that identifies the shock front at different times.

Regarding the wave properties, Figure D1 shows the pressure–volume (p – V) diagram for the shocked state and a quiescent condition for the event presented in Figure 4. For each snapshot ($t = 5812$ s to 5820 s), we extract the thermodynamic variables pressure, density, and temperature on the shock surface (p_1, ρ_1, T_1), while the preshock values (p_0, ρ_0, T_0) are extracted 2 s prior to the arrival of the shock on the same surface. For characterization of the quiescent condition, we use the thermodynamic variables at 2 and 4 s before the arrival of the shock at the surface. In any case, the simulation data take a realistic equation-of-state with variable adiabatic index into account. The variable V is the specific Volume $1/\rho$. The quiescent conditions, marked as gray dots in Figure D1, follow closely the black isentropic ($p V^\gamma = \text{constant}$) curve with minor volume compression/expansion typical of acoustic disturbances and temperature fluctuations that do not exceed 392 K. The shocked state (colored dots) still essentially follows the isentropic curve but deviates from the quiescent state in that it shows temperature changes of up to 2429 K with compression and a pressure enhancement of up to 3.5 times the preshock value. Yet, the preshock speeds relative to the shock front stay close to sonic, indicative of a weak shock (Y. B. Zel’dovich & Y. P. Raizer 1966). The shock wave

starts in a high β environment as a predominantly acoustic shock at $t \approx 5790$ s (see Figure 7) but continues as a slow mode, still predominantly acoustic, in the time frame 5812–5820 s. Therefore, we can compare it with results from pure hydrodynamics. The spread in the shocked state is due to the spread in the adiabatic index γ in the shocked media. In the case of a magnetic field-free, perfect gas with constant adiabatic index γ , the shocked state will lie on the Hugoniot curve given as

$$\frac{p_1}{p_0} = \left(\frac{\gamma + 1}{\gamma - 1} - \frac{V_1}{V_0} \right) / \left(\frac{\gamma + 1}{\gamma - 1} \frac{V_1}{V_0} - 1 \right) \quad (\text{D1})$$

from Equation (1.75) of Y. B. Zel’dovich & Y. P. Raizer (1966). The shock Hugoniot (red curve) closely follows the isentropic curve (black) for pressure changes of up to a factor 4 and starts deviating from the isentropic case for strong shocks only, which we do not have in the present situation. The close agreement of the numerical results (colored and gray dots) with the isentropic and the Hugoniot curve of an ideal, nonmagnetic fluid confirms the acoustic character of the wave and the difference in temperature and pressure fluctuations before and across the wave front its character as a weak shock. For a representation of the shock structure, Figure D2 shows the jump in temperature and gas pressure across the shock front along the white line of sight of Figure 4 at $(x, y) = (3435 \text{ km}, 3755 \text{ km})$ and time $t = 5800$ s. It shows the typical saw-tooth shape and the signature of a radiative precursor and relaxation of a radiative shock in a stratified




Figure D1. Pressure–volume (p – V) diagram showing the shocked and the quiescent condition. The temperature difference between pre- and postshock medium is displayed. The quiescent conditions (gray dots) follow the isentropic curve with minor volume compression and expansion typical of an acoustic disturbance. The temperature variations in the quiescent conditions are less than 392 K (marked by an arrow in the colorbar). The shocked state (colored dots) still essentially follows the isentropic curve but shows larger temperature changes with compression and pressure enhancements. Yet, plasma speeds relative to the shock front stay close to sonic, indicative of a weak shock. For comparison, the red and black curves show the Hugoniot and isentropic tracks, respectively, for a constant, mean $\gamma = 1.33$.




Figure D2. Temperature and gas pressure across the shock front. Temperature (red) and gas pressure (blue) along the white line of sight of Figure 4 at $(x, y) = (3435 \text{ km}, 3755 \text{ km})$ and time $t = 5800 \text{ s}$.

atmosphere as discussed in D. Mihalas & B. W. Mihalas (1984, Chapter 8.2) or by M. Carlsson & R. F. Stein (1997).

The wave energy flux is estimated on the constructed shock surface as $F = \Delta p v$, where Δp is the pressure perturbation relative to the background undisturbed medium 2 s prior to the arrival of the shock, and v is the velocity on the shock surface. The energy flux F is integrated over the area of the shock surface to estimate the power of the event quoted in Section 4.1.

We estimate the contribution to the chromospheric emission index $R'_{\text{HK}} = \log(F_{\text{HK}}/F_{\text{bol}})$, where F_{HK} is the radiative flux in the cores of the Ca II H and K spectral lines and $F_{\text{bol}} = \sigma T_{\text{eff}}^4$, the bolometric flux, where $\sigma = 5.7 \times 10^{-5} \text{ erg cm}^{-2} \text{ s}^{-1} \text{ K}^{-4}$ is the Stefan–Boltzmann constant and T_{eff} is the effective temperature, here 5778 K. For simulation event Nr. S1, we estimate that the shock front supplies a mean energy flux of $F_{\text{shock}} = 1.12 \pm 0.16 \times 10^7 \text{ erg cm}^{-2} \text{ s}^{-1}$ for about 8 s over a mean shock-front area of $A = 2.52 \pm 0.30 \times 10^{15} \text{ cm}^2$. Assuming that all this energy flux converts into H and K emission, and that the solar surface is at any time occupied with such shock surfaces with a filling factor f , we can compute $R'_{\text{HK}} = \log(f F_{\text{shock}}/F_{\text{bol}}) = \log f - 3.748$. The chromospheric basal flux of the Sun is $R'_{\text{HK}} = -5.2$ so that a filling factor of $\log f = -5.2 + 3.75 = -1.45$, hence $f = 0.035$, would suffice to supply the chromospheric basal flux by these shock events alone. Using the mean area A of shock event Nr. S1, this translates to one such event per $A/f = 7.2 \text{ Mm}^2$ or an occurrence rate of 0.14 per Mm^2 . Derivation of F_{HK} from the DKIST Ca II K filtergrams alone is conceivable but presently not possible. This is due to calibration issues, whereby the dimensional fluxes in $\text{erg str}^{-1} \text{ s}^{-1}$ are not available.

ORCID iDs

Friedrich Wöger <https://orcid.org/0000-0001-6907-9739>
 Gangadharan Vigeesh <https://orcid.org/0000-0002-9820-9114>
 Peter H. Keys <https://orcid.org/0000-0001-8556-470X>
 Oskar Steiner <https://orcid.org/0000-0003-4708-1074>
 Shahin Jafarzadeh <https://orcid.org/0000-0002-7711-5397>
 Robertus Erdélyi <https://orcid.org/0000-0003-3439-4127>
 David B. Jess <https://orcid.org/0000-0002-9155-8039>
 Mihalís Mathioudakis <https://orcid.org/0000-0002-7725-6296>
 Alexandra Tritschler <https://orcid.org/0000-0003-3147-8026>
 Damian J. Christian <https://orcid.org/0000-0003-1746-3020>
 Andre Fehlmann <https://orcid.org/0000-0002-7978-368X>
 Sarah Jaeggli <https://orcid.org/0000-0001-5459-2628>

References

- Battaglia, A. F., Canivete Cuissa, J. R., Calvo, F., Bossart, A. A., & Steiner, O. 2021, *A&A*, **649**, A121
- Beeck, B., Schüssler, M., Cameron, R. H., & Reiners, A. 2015, *A&A*, **581**, A42
- Bellot Rubio, L., & Orozco Suárez, D. 2019, *LRSP*, **16**, 1
- Berger, T. E., & Title, A. M. 1996, *ApJ*, **463**, 365
- Berger, T. E., & Title, A. M. 2001, *ApJ*, **553**, 449
- Berger, T. E., Rouppe van der Voort, L. H. M., Löfdahl, M. G., et al. 2004, *A&A*, **428**, 613
- Bloomfield, D. S., Lagg, A., & Solanki, S. K. 2007, *ApJ*, **671**, 1005
- Bonet, J. A., Márquez, I., Sánchez Almeida, J., Cabello, I., & Domingo, V. 2008, *ApJL*, **687**, L131
- Boro Saikia, S., Marvin, C. J., Jeffers, S. V., et al. 2018, *A&A*, **616**, A108
- Calvo, F. 2018, PhD thesis, Université de Genève
- Canivete Cuissa, J. R., Riva, F., & Steiner, O. 2024, *A&A*, **689**, A198
- Canivete Cuissa, J. R., & Steiner, O. 2024, *A&A*, **682**, A181

- Carlsson, M., De Pontieu, B., & Hansteen, V. H. 2019, *ARA&A*, **57**, 189
- Carlsson, M., Hansteen, V. H., Gudiksen, B. V., Leenaarts, J., & De Pontieu, B. 2016, *A&A*, **585**, A4
- Carlsson, M., & Stein, R. F. 1997, *ApJ*, **481**, 500
- Cherry, G., Gudiksen, B., & Finley, A. J. 2025, *A&A*, **700**, A276
- Cherry, G., Gudiksen, B., Finley, A. J., & Noraz, Q. 2026, *A&A*, **705**, A42
- Crockett, P. J., Mathioudakis, M., Jess, D. B., et al. 2010, *ApJL*, **722**, L188
- Dakanalis, I., Tsiropoula, G., Tziotziou, K., & Kontogiannis, I. 2022, *A&A*, **663**, A94
- De Pontieu, B., McIntosh, S. W., Carlsson, M., et al. 2007, *Sci*, **318**, 1574
- de Wijn, A. G., Stenflo, J. O., Solanki, S. K., & Tsuneta, S. 2009, *SSRv*, **144**, 275
- de Wijn, A. G., Casini, R., Carlile, A., et al. 2022, *SoPh*, **297**, 22
- Díaz-Castillo, S. M., Fischer, C. E., Rezaei, R., Steiner, O., & Berdyugina, S. 2024, *A&A*, **691**, A37
- Dumusque, X., Udry, S., Lovis, C., Santos, N. C., & Monteiro, M. J. P. F. G. 2011, *A&A*, **525**, A140
- Dunn, R. B., & Zirker, J. B. 1973, *SoPh*, **33**, 281
- Erdélyi, R., & Fedun, V. 2007, *Sci*, **318**, 1572
- Fedun, V., Shelyag, S., & Erdélyi, R. 2011, *ApJ*, **727**, 17
- Finley, A. J., Brun, A. S., Carlsson, M., et al. 2022, *A&A*, **665**, A118
- Fischer, C. E., Bello González, N., & Rezaei, R. 2017, *A&A*, **602**, L12
- Fischer, C. E., de Wijn, A. G., Centeno, R., Lites, B. W., & Keller, C. U. 2009, *A&A*, **504**, 583
- Fisher, G. H., & Welsch, B. T. 2008, *ASPC*, **383**, 373
- Fossum, A., & Carlsson, M. 2005, *Natur*, **435**, 919
- Freytag, B. 2013, *MSAIS*, **24**, 26
- Freytag, B., Steffen, M., Ludwig, H. G., et al. 2012, *JCoPh*, **231**, 919
- Gudiksen, B. V., Carlsson, M., Hansteen, V. H., et al. 2011, *A&A*, **531**, A154
- Hansteen, V. H., De Pontieu, B., Rouppe van der Voort, L., van Noort, M., & Carlsson, M. 2006, *ApJL*, **647**, L73
- Harten, A., Lax, P., & van Leer, B. 1983, *SIAMR*, **25**, 35
- Hasan, S. S., & Kalkofen, W. 1999, *ApJ*, **519**, 899
- Jafarzadeh, S., Solanki, S. K., Feller, A., et al. 2013, *A&A*, **549**, A116
- Jefferies, S. M., McIntosh, S. W., Armstrong, J. D., et al. 2006, *ApJL*, **648**, L151
- Jess, D. B., Jafarzadeh, S., Keys, P. H., et al. 2023, *LRSP*, **20**, 1
- Jess, D. B., Mathioudakis, M., Christian, D. J., et al. 2010, *SoPh*, **261**, 363
- Jess, D. B., Mathioudakis, M., Erdélyi, R., et al. 2009, *Sci*, **323**, 1582
- Keys, P. H., Mathioudakis, M., Jess, D. B., et al. 2011, *ApJL*, **740**, L40
- Keys, P. H., Reid, A., Mathioudakis, M., et al. 2019, *MNRAS*, **488**, L53
- Keys, P. H., Campbell, R. J., Magill, D. K. J., et al. 2026, *ApJ*, **999**, 201
- Khomenko, E., & Cally, P. S. 2019, *ApJ*, **883**, 179
- Kitiashvili, I. N., Kosovichev, A. G., Mansour, N. N., & Wray, A. A. 2012, *ApJL*, **751**, L21
- Lakeland, B. S., Naylor, T., Haywood, R. D., et al. 2024, *MNRAS*, **527**, 7681
- Liu, J., Carlsson, M., Nelson, C. J., & Erdélyi, R. 2019a, *A&A*, **632**, A97
- Liu, J., Nelson, C. J., Snow, B., Wang, Y., & Erdélyi, R. 2019b, *NatCo*, **10**, 3504
- Mihalas, D., & Mihalas, B. W. 1984, *Foundations of Radiation Hydrodynamics* (Oxford Univ. Press)
- Moll, R., Cameron, R. H., & Schüssler, M. 2012, *A&A*, **541**, A68
- Nagata, S., Tsuneta, S., Suematsu, Y., et al. 2008, *ApJL*, **677**, L145
- Noraz, Q., Carlsson, M., & Aulanier, G. 2026, *A&A*, **705**, A86
- Osborne, C. M. J., & Milić, I. 2021, *ApJ*, **917**, 14
- Park, S. H., Tsiropoula, G., Kontogiannis, I., et al. 2016, *A&A*, **586**, A25
- Pereira, T. M. D., & Uitenbroek, H. 2015, *A&A*, **574**, A3
- Pérez Martínez, M. I., Schröder, K.-P., & Cuntz, M. 2011, *MNRAS*, **414**, 418
- Priest, E. 2014, *Magnetohydrodynamics of the Sun* (Cambridge Univ. Press)
- Rimmele, T. R., Warner, M., Keil, S. L., et al. 2020, *SoPh*, **295**, 172
- Riva, F., & Steiner, O. 2022, *A&A*, **660**, A115
- Rutten, R. J., Kiselman, D., Rouppe van der Voort, L., & Plez, B. 2001, *ASPC*, **236**, 445
- Salhab, R. G., Steiner, O., Berdyugina, S. V., et al. 2018, *A&A*, **614**, A78
- Scalisi, J., Oxley, W., Ruderman, M. S., & Erdélyi, R. 2021, *ApJ*, **911**, 39
- Scullion, E., Erdélyi, R., Fedun, V., & Doyle, J. G. 2011, *ApJ*, **743**, 14
- Silva, S. S. A., Verth, G., Rempel, E. L., et al. 2021, *ApJ*, **915**, 24
- Spruit, H. C. 1976, *SoPh*, **50**, 269
- Spruit, H. C. 1979, *SoPh*, **61**, 363
- Stefani, F., Forbriger, J., Gundrum, T., Herrmannsdörfer, T., & Wosnitza, J. 2021, *PhRvL*, **127**, 275001
- Steiner, O., Grossmann-Doerth, U., Knölker, M., & Schüssler, M. 1998, *ApJ*, **495**, 468
- Steiner, O., Hauschildt, P. H., & Bruls, J. 2001, *A&A*, **372**, L13
- Title, A. M., & Berger, T. E. 1996, *ApJ*, **463**, 797
- Tsuneta, S., Ichimoto, K., Katsukawa, Y., et al. 2008, *ApJ*, **688**, 1374
- Tziotziou, K., Scullion, E., Shelyag, S., et al. 2023, *SSRv*, **219**, 1
- Uitenbroek, H. 2001, *ApJ*, **557**, 389
- Ulmschneider, P., Zähringer, K., & Musielak, Z. E. 1991, *A&A*, **241**, 625
- Vigeesh, G., Fedun, V., Hasan, S. S., & Erdélyi, R. 2012, *ApJ*, **755**, 18
- Wedemeyer-Böhm, S., & Rouppe van der Voort, L. 2009, *A&A*, **507**, L9
- Wedemeyer-Böhm, S., Scullion, E., Steiner, O., et al. 2012, *Natur*, **486**, 505
- Wöger, F. 2014, *SPIE*, **9147**, 914791
- Wöger, F., Rimmele, T., Ferayorni, A., et al. 2021, *SoPh*, **296**, 145
- Wöger, F., von der Lühe, O., & Reardon, K. 2008, *A&A*, **488**, 375
- Zel'dovich, Y. B., & Raizer, Y. P. 1966, in *Physics of Shock Waves and High-Temperature Hydrodynamic Phenomena*, ed. D. Wallace & R. Probstein, Vol. I (Academic Press)